

Although you are outnumbered,
together we can fight the bacteria
that cause periodontal disease.



FACE IT.
YOU'RE OUTNUMBERED.

ARESTIN®—along with SRP—starts working immediately, fights infection and inflammation for 30 days and provides significantly better results than SRP alone for up to 90 days.^{1,2,3}

For full prescribing information or to learn more about ARESTIN®, ask your dental professional or visit www.arestin.com

ARESTIN® (minocycline hydrochloride) Microspheres, 1 mg is indicated as an adjunct to scaling and root planing (SRP) procedures for reduction of pocket depth in patients with adult periodontitis. ARESTIN® may be used as part of a periodontal maintenance program which includes good oral hygiene, and scaling and root planing. ARESTIN® contains minocycline, a tetracycline derivative, and therefore should not be used in children and in pregnant or nursing women. The use of drugs of the tetracycline class during tooth development may cause permanent discoloration of the teeth. The most common treatment-emergent adverse events were headache (9.0%), infection (7.6%), flu syndrome (5.0%), and pain (4.3%). These occurred at a similar rate to SRP and SRP + placebo.

References: **1.** Goodson JM, Gunsolley JC, Grossi SG, et al. Minocycline HCl microspheres reduce red-complex bacteria in periodontal disease therapy. *J Periodontol* 2007;78(8):1568-1579. **2.** Oringer RJ, Al-Shammari KF, Aldredge WA, et al. Effect of locally administered minocycline microspheres on markers of bone resorption. *J Periodontol* 2002;73:835-842. **3.** Williams RC, Paquette DW, Offenbacher S, et al. Treatment of periodontitis by local administration of minocycline microspheres: a controlled trial. *J Periodontol* 2001;72:1535-1544.

©2009 OraPharma, Inc. A-836-09 4/09

ARESTIN® is a registered trademark of OraPharma, Inc.

LISTERINE® Antiseptic and BRUSH, FLOSS & RINSE™ are trademarks of Johnson & Johnson Healthcare Products Division of McNEIL-PPC, Inc.


PERIODONTAL DISEASE MEANS YOU HAVE
MILLIONS OF HARMFUL BACTERIA CAUSING
DAMAGE DEEP UNDER YOUR GUMLINE—WHERE
BRUSHING AND FLOSSING CAN'T REACH.

Give your gums a fighting chance
against this infection with ARESTIN®
and professional treatment.

Arestin® 
minocycline HCl 1mg
MICROSPHERES

Periodontal disease is a serious infection under your gumline.

ARESTIN® + SCALING & ROOT PLANING (SRP) KILL BACTERIA THAT CAUSE PERIODONTAL INFECTIONS.



ARESTIN® is an antibiotic placed directly in the infected areas—or “pockets”—in your gums. It’s applied right after scaling and root planing, the dental procedure that disrupts stubborn plaque and bacteria below your gumline.

ARESTIN® FIGHTS INFECTION AND INFLAMMATION FOR 30 DAYS AND PROVIDES BETTER RESULTS THAN SRP ALONE FOR UP TO 90 DAYS.^{1,2,3}

The ARESTIN® microspheres gradually release minocycline, an antibiotic which kills bacteria that cause periodontal infection, inflammation, and bleeding.

ARESTIN® starts working quickly and keeps fighting bacteria long after you leave the dental office.²

INSIDE YOUR MOUTH

DATE: _____



NOTES:

UNDERSTANDING WHAT YOUR NUMBERS MEAN.



A pocket of 4mm or more in size may mean you have an infection.

After Your ARESTIN® Treatment

Here are a few simple steps to help you get the maximum results from your treatment.

- Avoid eating hard, crunchy, or sticky foods for **one week**.
- Don't touch treated areas for **one week**.
- Wait **10 days** to floss around treated areas.

Remember:

- Practice good oral hygiene by sticking to a daily BRUSH, FLOSS & RINSE™ routine, using an ADA-accepted antimicrobial rinse such as LISTERINE® Antiseptic. Use as directed.
- Follow up with your dentist and your dental hygienist by keeping your scheduled appointments.